INDEX TO VOLUME

New names an d the final members of new combinations are in **heavier type.**

Abies grandis, 99; Mertensiana, 100; Pattoniana, 100
Acalypha **jamaicensis,** 7; pruinosa, 7
Acer platanoides, 408; rubrum, 407; saccharinum, 408
Aconiopteris gorgonea, 572; obtusa, 572
Acorus Calamus, 107
Acrophorus Macraeanus, 582; repens, 582
Acrostichum, 258, 571; aemulum, 571, 572; calomelanos, 588; crassifolium, 572; gorgonium, 572; Helleri, 571; hirtum, 570; lineatum, 275; micans, 570; micradenium, 570; pellucido-marginatum, 572; reticulatum, 572; Spicant, 275; squamosum, 570; vestitum, 570; Wawrae, 571
Actinostemon **jamaicensis,** 7
Adiantum, 568, 577; Bennettii, 577; Capillus, 577; Capillus-veneris, 577; chinense, 585; chusanum, 585; emarginatum, 577; Levingei, 577; tenuifolium, 585; Wattii, 577.
Adnaria, 550
Aenothera grandiflora, 519, 521; longiflora, 521
Ailanthus glandulosa, 420
ALDEN, I. A contribution to the life history of Uvularia sessilifolia, 439
Alectoria jubata, 201; pubescens, 201
Alisma brevipes, 101; Geyeri, 101; Plantago-aquatica, 101
Alismaceae, 101
Alliaceae, 110
Allium, 16, 19, 22; Cepa, 15, 17, 20
Allosorus aquilinus, 575; ternifolius, 576
Alnus, 439
Alopecurus aristulatus, 102; fulvus, 102; pallescens, 102
Alsinaceae, 314
Alsine **alpestris,** 315; Edwardsii, 315; **Palmeri,** 315; stricta, 315; **strictiflora,** 315; **subvestita,** 315
Alsinopsis **dawsonensis,** 316; **pusilla,** 316
Amaranthaceae, 313
Amaranthus californicus, 313; carneus, 313; graecizans pubescens, 313; **pubescens,** 313
Ammodenia **oblongifolia,** 317
Ampelocissus Alexandri, 12
Ampelopsis, 39, 50
Anacardiaceae, 423
ANDREWS, F. M. Protoplasmic streaming in Mucor, 455
Andromeda Novae-caesareae, 390, 405; Pfaffiana, 388; Snowii, 390, 405
Androvettia, 389
Anetium reticulatum, 572
Anticlea, 108; coloradensis, 108; elegans, 108; porrifolia, 108, 109; **vaginata,** 108, 109
Apios Apios, 80
Aphanolejeunea, 222, 224, 329
Arabis **brevisiliqua,** 326; lignifera, 326; **MacDougalii,** 326; subpinnatifida, 326
Araceae, 107
Aralia cottondalensis, 403; decurrens, 404; Saportana, 403, 404; Wellingtoniana, 390, 402–404; Wellingtoniana Vaughanii, 388
Araucaria bladenensis, 389; Jeffreyi, 389
Arenaria **cephaloidea,** 316; congesta, 316; dawsonensis, 316; diffusa, 316; Fendleri diffusa, 316; laricifolia, 316; **laxiflora,** 316; **lithophila,** 316; peploides major, 317; pusilla, 316; sitchensis, 317; subcongesta lithophila, 316
Aristida bromoides, 102; fasciculata Hookeri, 102; oligantha, 102; longiseta, 102
Arrhenatherum elatius, 103
Artabotrys odoratissima, 502
Arthrotaxopsis, 393
Ascobolaceae, Methods of culture and the morphology of the archicarp in certain species of the, 139
Ascobolus, 140, 141, 144–146, 149, 150, 167, 170, 171, 187, 188; aerugineus, 151; albidus, 144, 148, 149; amoenus, 179; atrofuscus, 171; brunneus, 179; carbonarius, 141, 148, 149, 151, 159–163, 170–172, 187, 189–195; carbonicola, 171; denudatus, 143, 149; furfuraceus, 141, 144–149, 151, 168, 169, 185, 186, 192–194; glaber, 146, 151, 168, 169, 180, 193, 194; immersus, 142, 149, 151, 164, 165, 169, 184, 186, 190, 193, 194; Leveillei, 151, 166, 169, 179, 180, 194; Leveillei americanus, 152, 167, 169, 179; mirabilis, 191; perplexans, 144, 148; pulcherrimus, 457; pusillus, 151, 152, 170, 172, 193; sp., 148, 151, 156; stictoideus, 179; viridis, 141, 148, 151, 152, 164, 171, 172, 193; viridulus, 152, 170; Winteri, 151, 152, 154, 156, 158, 179, 180, 181, 184, 186, 190, 193, 195; xylophilus, 167, 169, 193

Ascodesmis, 139, 186, 191; nigricans, 141, 145, 148, 152, 169, 170
Ascophanus, 187; Aurora, 152; carneus, 144, 147–149, 151, 152, 169, 178–180, 186, 190, 193, 457, 491; glaucellus, 152; granuliformis, 152; Holmskjoldii, 152; lacteus, 152; microsporus, 152; minutissimus, 152; ochraceus, 146, 152, 190; pilosus, 142; saccharinus, 143, 148; sarcobius, 152, 169, 178, 193
Asparagin, effect of, on absorption and growth in wheat, 429
Aspergillus, 149, 190; niger, 535
Aspidium cordifolium, 583; cyatheoides, 599; Ecklonii, 598; exaltatum, 584; Filix-mas fusco-atrum, 592; glabrum, 592; glabrum pusillum, 593; glabrum quadripinnatum, 592; globuliferum, 591; hawaiiense, 594; Hudsonianum, 601; juglandifolium macrosorum, 26; latifrons, 593; molle, 599; nymphale, 599; paleaceum, 591; parallelogrammum, 591; parasiticum, 599; patens, 599; propinquum, 598; resiniferum, 598; rubiginosum, 594; Schkuhrii, 584; Speluncae, 586; squamigerum, 594; tuberosum, 583; undulatum, 583; unitum, 598; violascens, 599
Asplenium, 258, 569; aquilinum, 575; Spicant, 275
Athyrium, 569
Atriplex aptera, 311; **buxifolia,** 311; canescens, 311, 312; confertifolia, 312; **Garretti,** 312; Nuttallii, 311; occidentalis, 312; odontophora, 311; phyllostegia, 312; **tetraptera,** 311, 312
Avena americana, 103; Hookeriana, 103; Mortoniana, 103
Azolla, 232

Baptisia tinctoria, 69
Batodendron, 549, 553
BEATTIE, J. H., and SKINNER, J. J. Effect of asparagin on absorption and growth in wheat, 429
Benzoin venustum, 388, 390, 399
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII. Texas, 387; Notes on the genus Widdringtonites, 341
BICKNELL, E. P. The ferns and flowering plants of Nantucket—IX, 69; —X, 415
Bicornia, 553
Bidens **clarendonensis,** 9
Blechnum, 257–259, 261, 276; attenuatum, 272; boreale, 275; capense, 367, 375; Christii, 362; costaricensis, 363; danaeaceum, 364; doodioides, 275, 277; ensiforme, 263; falciforme, 365, Ghiesbreghtii, 277; L'Herminieri, 268; lineatum, 367; onocleoides, 271; ornifolium, 371; **Plumieri,** 269; polypodioides, 272; sessilifoli[um], 373; Spicant, 275; stoloniferum, 277; striatum, 375; tubulare, 377; varians, 379; violaceum, 379; Werckleanum, 382
Blepharidachne Kingii, 103
Boryana [Struthiopteris], 378
Botrychium, 234; subbifoliatum, 234
Boudiera, 152; Claussenii, 152, 170
Brachyphyllum crassicaule, 393; macrocarpum **formosum,** 389, 390, 392, 406; obesiforme elongatum, 393
BRAINERD, E. Violet hybrids between species of the palmata group, 85
Brassicaceae, 322
BRITTON, N. L. Studies of West Indian plants—IV, 1
Briza maxima, 104
BROADHURST, J. The genus Struthiopteris and its representatives in North America, 257, 357
Brodiaea, 110, 111; capitata, 111; capitata pauciflora, 111; congesta, 111; grandiflora, 110, 111; insularis, 111; volubilis, 111

Caeoma nitens, 113–116
Caesalpiniaceae, 69
Calamagrostis, 103; lucida, 103; rubescens, 103; Vaseyi, 103
Callitrichaceae, 422
Callitriche heterophylla, 422
Callitris, 342, 343
Calyptridium roseum, 314
Canangium **Brandesanum,** 504; odoratum, 504
Cameraria angustifolia, 5; latifolia, 5, 6; **microphylla,** 6; **oblongifolia,** 6; retusa, 5
Caryophyllaceae, 317
Case of changed polarity in Spirogyra elongata, A, 509
Cassia Chamaechrista, 69
Cathartolinum intercursum, 418
Celtis Douglasii, 305; occidentalis, 305; reticulata, 304, 304; **rugosa,** 304
Cerastium arvense fuegianum, 314, 315; fuegianum, 314; strictum, 315; **thermale,** 314
Ceropteris, 568, 588; calomelanos, 588; **ochracea,** 588
Cetraria nivalis, 201; islandica, 201
Chara, 467
Cheilanthes, 579; Lidgatii, 579; ternifolia, 576
Cheilolejeunea decidua, 279
Cheiranthus, 323; argillosus, 324; aridus, 323, 324; asper, 323; asperrimus, 324; Bakeri, 324; cheiranthoides, 323; elatus, 323; inconspicuus, 323; nivalis, 324; nivalis amoenus, 324; Pallasii, 324; pygmaeus, 324; syrticola, 323; Wheeleri, 324
Cheirinia, 323; **amoena,** 324; **argillosa,**

324; **arida,** 323; **aspera,** 323, 325; **asperrima,** 324; **Bakeri,** 324; **brachycarpa,** 325; cheiranthoides, 323; **elata,** 323; **inconspicua,** 323; **nivalis,** 324; **oblanceolata,** 324, 325; **Pallasii,** 324; **radicata,** 324; **syrticola,** 323; **Wheeleri,** 324
Chenopodiaceae, 310
Chenopodium calceoliforme, 313; desiccatum, 311; lanceolatum, 311; leptophyllum, 310; oblongifolium, 310; opulifolium, 311; paganum, 311; **praetericola,** 310; rubrum, 311; succosum, 311; viride, 311
Chiogenes, 550
Chloris brevispica, 103
Chorizanthe brevicornu, 310; **spathulata,** 309
Cibotium, 242; Chamissoi, 242, 243; glaucum, 243, 244; Menziesii, 242, 243; pruinatum, 243
Cinna arundinacea, 103
Cinnamomum, 400; ellipsoideum, 388; Heerii, 388; membranaceum, 390, 401; sp., 388
Cincinalis aquilina, 575
Cissus, 39, 41, 50
Cladophora, 509
Clerodendron **calcicola,** 9; spinosum, 10
Clusia **clarendonensis,** 7
Cocculus, 400
Codorolla, 554
Cololejeunea, 210–212, 222–234, 329; Biddlecomiae, 223; floccosa, 607; peraffinis, 608; platyneura, 608
Colura, 209, 210
Colurolejeunea, 329
Colutea primordialis, 390, 396
Commelinaceae, 107
Coniogramme, 568, 569; fraxinea, 589
Contribution to the life history of Uvularia sessilifolia, A, 439
Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII. Texas, 387
Coprinus, 186
Corema Conradii, 422
Cornophyllum vetustum, 390, 404
Coronaria, 317
Coronilla varia, 74
Corylus, 439
Cracca virginiana, 74
Cuba, Undescribed species from, 9
Cuba, The genus Ginoria in, 12
Cubonia sp., 152
Cunninghamites, 389
Cyatheaceae, 237, 241
Cyclolejeunea, 329
Cymbopetalum penduliflorum, 502
Cyrtomium, 569
Cystopteris Douglasii, 587; sandwichensis, 587
Cystopus, 188
Cytisus scoparius, 70

DACHNOWSKI, A. The relation of Ohio bog vegetation to the chemical nature of peat soils, 53
Danthonia spicata, 103
DARLING, C. A. Mitosis in living cells, 407
Davallia biflora, 584; chinensis, 585; chusana, 585; flaccida, 586; Khasiyana, 586; Macraeana, 582; Mannii, 582; polypodioides, 585; remota, 585; Speluncae, 586; strigosa, 586; tenuifolia, 585
DAVIS, B. M. Was Lamarck's evening primrose (Oenothera Lamarckiana Seringe) a form of Oenothera grandiflora Solander?, 519
Delphinium bicolor, 320, 322; **coelestinum,** 320; Geyeri, 322; **Helleri,** 321; **Leonardi,** 320; multiflorum, 322; reticulatum, 322; scaposum, 321; scopulorum, 322; **viscidum,** 322; **xylorrhizum,** 321
Dendropanax arboreum, 1, 2; **blakeanum,** 4; **brachypodum,** 2; **cordifolium,** 4; cuneifolium, 2; **elongatum,** 3; **grande,** 4; **grandiflorum,** 3; laurifolium, 2; nutans, 2; pendulum, 2, 4; samydifolium, 1
Dendropanax in the West Indies, 1
Dennstaedtia strigosa, 586
Deschampsia caespitosa, 103; **pungens,** 103
Desmos, 504, 505; chinensis, 502, 504; cochinchinensis, 504–506; **crinitus,** 507; **dasymaschalus,** 507; **desmanthus,** 508; **dumosus,** 506; **Dunalii,** 506; **elegans,** 506; **Lawii,** 506; **longiflorus,** 507; **pannosus,** 506; **praecox,** 507; **stenopetalus,** 507; **viridiflorus,** 506; **Wrayi,** 507; **zeylanicus,** 506
Desmos the proper generic name for the so-called Unonas of the Old World, 501
Detonia trachycarpa, 170
Diastoloba, 280
Dichasium parallelogrammum, 591
Dichelostemma, 110, 111; congesta, 111; multiflora, 111
Dicksonia glauca, 244; japonica, 585; Menziesii, 243; splendens, 243; strigosa, 585
Dicranopteris, 238, 239; dichotoma, 241; **emarginata,** 239, 240; glabra, 239, 240; glauca, 239, 240; linearis, 239, 240; **owhyhensis,** 239, 241
Didymoglossum humile, 246
Diellia, 568, 580; Alexandri, 580, 581; **centifolia,** 580; erecta, 580, 581; falcata, 580, 581; Knudsenii, 580, 582; **laciniata,** 580, 581; **Mannii,** 580, 582; pumila, 580
Diospyros primaeva, 388; Steenstrupi, 388

Diplasiolejeunea, 209, 210, 212, 222, 224, 329, 603, 605; armatiloba, 216; **brachyclada,** 216, 217, 220, 222, 225, 605; **Johnsonii,** 603–606, 611; pellucida, 210–225, 605, 606; pellucida **malleiformis,** 215, 222; Rudolphiana, 220–222, 224, 225, 605; unidentata, 217, 219, 221, 222, 224, 225, 605
Diplasiolejeunea, Vegetative reproduction in, 222
Diplazium fraxineum, 589
Diplusodon ginorioides, 13
Dipterostemon, 110; **capitatus,** 111; **insularis,** 111; **pauciflorus,** 111; **pulchellus,** 111
Discoid gemmae in Radula, 329
Disterigma, 552
DODGE, B. O. Methods of culture and the morphology of the archicarp in certain species of the Ascobolaceae, 139
Dondia **calceoliformis,** 313
Doodia, 276, 569
DORSEY, M. J. Variation in the floral structures of Vitis, 37
Doryopteris, 567, 576; decipiens, 576; decora, 576, 577; pedata, 576
Draba andina, 327, 328; aurea, 328; aureformis, 328; decumbens, 328; deflexa, 328; densiflora, 327, 328; glacialis pectinata, 327; lapilutea, 328; luteola, 328; **pectinata,** 327, 328; praealta, 328; reflexa, 328; surculifera, 328; uber, 328; yellowstonensis, 328
Dryopteris, 568, 569, 589; acutidens, 590, 597; crinalis, 590, 595; cyatheoides, 591, 599–601; dilatata, 597; fijiensis, 595; Filix-mas, 592, 594; **fusco-atra,** 589, 592; glabra, 590, 592, 594; globulifera, 589, 591, 599; **hawaiiensis,** 590, 594; honolulensis, 590, 595; Keraudreniana, 590, 596; latifrons, 590, 593, 595; nuda, 590, 592, 593; **paleacea,** 589, 591, 592; parasitica, 591, 598, 599; **parvula,** 590, 593; propinqua, 590, 598; pusilla, 593; **rubiformis,** 590, 596; rubiginosa, 590, 594; sandwicensis, 590, 598, 600; spinulosa, 592, 597; squamigera, 590, 594; stegnogrammoides, 591, 600, 601; truncata, 591, 601; unidentata, 590, 597
Effect of asparagin on absorption and growth in wheat, 429
Effect of guanidin on plants, 535
Eichhornia, 16, 19, 21, 22; speciosa, 15
Elaeagia **cubensis,** 10
Elaphoglossum, 567, 569, 571; aemulum, 570, 571; conforme, 569; gorgonium, 570, 572; hirtum, 570; micradenium, 570; nitidum, 570; squamosum, 570; reticulatum, 570, 572; vestitum, 570; Wawrae, 570, 571
Elisanthe, 318; Drummondii, 318
Elymus, 107
Empetraceae, 422
Endolepis **phyllostegia,** 312
Ephedra nevadensis, 100; viridis, 100
Ephedraceae, 100
Eragrostis hypnoides, 104; lutescens, 104; neomexicana, 104; secundiflora, 104
Eremochloa Kingii, 103
Ericacea, 553
Ericales, 405
Eriocoma cuspidata, 102; **hymenoides,** 102
Eriogonum, 308, 309; androsaceum, 309; angulosum, 309; aureum, 308, 309; Baileyi, 309; bicolor, 309; **biumbellatum,** 306; campanulatum, 307; chloranthum, 308; clavellatum, 309; compositum, 309; corymbosum, 308; crassifolium, 308; croceum, 307; deflexum, 309; densum, 309; depauperatum, 309; **depressum,** 308; dichotomum, 309; elatum, 309; flavum, 308; Hookeri, 309; **idahoense,** 307; insigne, 309; leptophyllum, 309; longilobum, 309; Mearnsii, 309; micranthum, 309; microthecum, 307; neglectum, 307; nidularium, 309; nutans, 309; ochrocephalum, 309; Ordii, 309; orendense, 307; ovalifolium depressum, 308; Parryi, 309; polifolium, 309; Porteri, 309; proliferum, 309; puberulum, 309; pulvinatum, 309; racemosum, 309; ramosissimum, 308, 309; rubidum frigidum, 308; salicinum, 308; Shockleyi, 309; spathulatum, 308, 309; **spathuliforme,** 307; spergulinum, 309; strictum, 308, 309; subreniforme, 309; sulcatum, 309; tenellum, 309; Thomasii, 309; Thompsonae, 309; thymoides, 309; trichopodum, 309; tristichum, 308; turbinatum, 309; umbellatum, 307; umbelliferum, 307; villiflorum, 309; vimineum, 309; Wetherillii, 309; Wrightii, 309
Erodium cicutarium, 417
Erysimum, 323; alpestre, 324; amoenum, 324; argillosum, 324; asperum, 323; asperum alpestre, 324; asperum inconspicuum, 323; Bakeri, 324; cheiranthoides, 323; elatum, 323; inconspicuum, 323; nivale, 324; oblanceolatum, 324; officinale, 323; parviflorum, 323; pygmaeum, 325; radicatum, 324; syrticolum, 323; Wheeleri, 324
Erythronium, 441
ESTABROOK, A. H., and LIVINGSTON, B. E. Observations on the degree of stomatal movement in certain plants, 15
Eudianthe, 317
Eucallitris, 342
Eucalyptus Geinitzi, 390, 402
Eugenia primaeva, 388
Euphorbia Cyparissias, 422; maculata, 421; polygonifolia, 421; Preslii, 421

Euphorbiaceae, 421
Eupteris aquilina, 575
Eurotia lanata, 312; **subspinosa,** 312, 313
EVANS, A. W. Hepaticae of Puerto Rico — XI. Displasiolejeunea, 209; New West Indian Lejeuneae—II, 603

Falcata monoica, 80; Pitcheri, 80
Ferns and flowering plants of Nantucket, The—IX, 69; —X, 415
Ferns from Tropical America, New, 285
Festuca, 105, 106; campestris, 105; confinis, 106; dasyclada, 105; Kingii, 106; megalura, 105; ovina calligera, 105; pacifica, 105; reflexa, 105; scabrella, 105; Thurberi, 105; viridula, 105; Watsonii, 106
Ficus daphnogenoides, 390, 394; glascoeana, 388
Filix, 568, 587; **Douglasii,** 587
Filicales, 232, 236
Flax rust, 113
Frenela, 342
Frenelites, 343
Frenelopsis, 343, 346, 347; bohemica, 346; Heheneggeri, 346; ramosissima, 346, 347
FROMME, F. D. Sexual fusions and spore development of the flax rust, 113
Frullania, 279, 280, 284, 329; Asagrayana, 280; Bolanderi, 279, 284; Brittoniae, 280, 283, 284; cleistostoma, 284; dilatata, 280, 283; eboracensis, 279, 280, 281, 283, 284; fragilifolia, 280; inflata, 280; plana, 280, 283, 284; riparia, 280, 283, 284; saxicola, 280; Selwyniana, 280; squarrosa, 221; Tamarisci, 280
Frullaniae, Vegetative reproduction in the New England, 279
Funkia, 19, 22; ovata, 15

Gaultheria, 550
Gaylussacia, 549, 550; buxifolia, 550; dumosa, 551; frondosa, 555; nana, 555; tomentosa, 555
Genus Cameraria, The, 5
Genus Ginoria in Cuba, The, 12
Genus Struthiopteris and its representatives in North America, The, 257, 357
Geraniaceae, 415
Geranium carolinianum, 415; maculatum, 415; molle, 416; pusillum, 416; Robertianum, 415
Gilibertia brachypoda, 2
Ginoria americana, 12, 13; **arborea,** 13; curvispina, 13; Diplusodon, 13; **ginorioides,** 13; spinosa, 13
Gleditsia triacanthos, 69
Gleichenia glauca, 240; hawaiiensis, 241; linearis, 241; longissima, 240; owhyhensis, 241
Gleicheniaceae, 237, 238
Gonocormus saxifragoides, 245
Guanidin, the effect of, on plants, 535
Guatteria, 502
Gymnoascus, 186; ruber, 186
Gymnoconia interstitialis, 113, 114, 116
Gymnogramme calomelanos, 588; javanica, 589; pilosa, 589; tartarea, 588
Gymnospermae, 391
Gymnosporangium clavariaeforme, 119, 128
Gymnotheca Douglasii, 236

Hawaiian Islands, A taxonomic study of the Pteridophyta of the, 227, 567
Hedera nutans, 3, 4,
Hepatica, 439
Hepaticae of Puerto Rico—XI. Diplasiolejeunea, 209
Hesperis pygmaeus, 325
Hesperochloa, 106; **Kingii,** 106
Hesperopeuce **Mertensiana,** 100; Pattoniana, 100
Hexaclinis, 342
Hicriopteris speciosa, 240
Homotropantha, 280
Honckenya oblongifolia, 317
Hookera, 111; coronaria, 110; pulchella, 111
Hordeum Aegiceras, 106; caespitosum, 106; montanense, 106; murinum, 106
HOWE, R. H. JR. The lichens of the Linnean Herbarium with remarks on Acharian material, 199
Hugeria, 551
Humaria, 190, 191; granulata, 191, 195
Humata Mannii, 582
Hymenodium crassifolium, 572; reticulatum, 572
Hymenophyllum, 244, 245, 247; Baldwinii, 247; lanceolatum, 247, 248; obtusum, 247, 248; recurvum, 247; tunbridgense, 247
Hymenophyllaceae, 237, 244
Hypocopra, 190
Hypolepis, 568, 578; **flaccida,** 578, 579; punctata, 578; tenuifolia, 578, 579

Ilex, 387; bronxensis, 426, 427; **fastigiata,** 426; glabra, 425; verticillata, 426, 427; verticillata padifolia, 426
Ilicaceae, 425
Index to American Botanical Literature, 29, 81, 133, 205, 249, 293, 349, 411, 447, 511, 561, 613
Inga cretacea, 388
Isatis, 20, 22; tinctoria, 15

Jamaica, Undescribed species from, 7
Juncaceae, 110
Juncus arizonicus, 110; columbianus, 110; ensifolius, 110; Jonesii, 110; mexi-

canus, 110; Regelii, 110; saximontanus, 110; Tracyi, 110; uncialis, 110
Jungermannia floccosa, 607; pellucida, 213, 214; unidentata, 217, 219
Juniperaceae, 99
Juniperites baccifera, 342
Juniperus, 281; horizontalis, 100; Knightii, 99; monosperma, 99; prostrata, 100; Sabina procumbens, 100; utahensis, 99, 100; virginiana, 99

Lachnea melaloma, 170; stercorea, 147, 190
Larix decidua, 408
Lasiobolus, 187; equinus, 142, 152, 168, 169; pulcherrimus, 140
Lastraea glabra, 592; globulifera, 591; latifrons, 593; rubiginosa, 594; squamigera, 594; truncata, 591
Lathyrus maritimus, 79; pilosus, 79; tuberosus, 80
Laurophyllum minus, 390, 402
Laurus, 387; plutonia, 390, 401; proteaefolia, 388
Leiolejeunea grandiflora, 603
Lejeunea, 210, 329, 603; albifolia, 214; floccosa, 607; ocellulata, 213, 214; pellucida, 213; platyneura, 608; unidentata, 217, 220
Lemna cyclostasa, 107; minima, 107; minor, 107; perpusilla, 107
Lemnaceae, 107
Lepidium **hirsutum,** 322; intermedium pubescens, 322; medium pubescens, 322; virginicum texanum pubescens, 322
Leptocolea, 210, 222–224, 329, 603, 607; **appressa,** 606–608; **floccosa,** 607, 608; scabriflora, 607
Lespedeza, 75; Bicknellii, 77; Brittonii, 76; capitata, 78; frutescens, 77; hirta, 77; Nuttallii, 76; procumbens, 75; Stuvei, 76; velutina, 77; violacea, 77; virginica, 76
Letharia vulpina, 201
Lichens of the Linnean Herbarium with remarks on Acharian material, The, 199
Limnia depressa, 314; spathulata, 314; **utahensis,** 314
Linaceae, 418
Lindera venusta, 399
Lindsaya Alexandri, 581; centifolia, 580; chinensis, 585; Knudsenii, 582; laciniata, 581; laciniata subbipinnata, 581; Mannii, 582; repens, 583
Linum, 419; floridanum, 419, 420; **intercursum,** 418–420; medium, 419; striatum, 418; sulcatum, 420; usitatissimum, 118, 418; virginianum, 418
Liquidambar integrifolium, 388
Liriodendron pinnatifidum, 388, 395; quercifolium, 390, 395; Snowii, 395
Litobrochia decora, 577; decipiens, 576
Litsea falcifolia, 399
LIVINGSTON, B. E., and ESTABROOK, A. H. Observations on the degree of stomatal movement in certain plants, 15
Lomaria, 257, 259; acrodonta, 371, 373; attenuata, 272, 383; Boryana, 377; brasiliensis, 375, 377; costaricensis, 363; crenata, 275, 277; danaeacea, 364, 365; decrescens, 271, 273; deflexa, 365; divergens, 271; ensiformis, 263; exaltata, 264, 266, 271; falciformis, 365; Féei, 272; fragilis, 271, 273; Gheisbreghtii, 277; L'Herminieri, 266, 267; lineata, 366; longifolia, 371, 372; martinicensis, 269; mexicana, 271, 273; mucronata, 266, 271; onocleoides, 271; ornifolia, 373; Plumieri, 269; polypodioides, 271; procera, 366, 383, 384; pteropus, 274; robusta, 369, 370; rufa, 369, 383; Ryani, 375, 377, 383; Schiedeana, 370–373; Schomburgkii, 383; sessilifolia, 373; spectabilis, 371, 372, 38 ; Spicant, 275; Spicant serra 377; spissa, 362; stolonifera, 277; striata, 375; tuberculata, 375; varians, 379; violacea, 379; Werckleana, 382
Lomaridium Herminieri, 267; Plumieri, 269
Lonchitis tenuifolia, 578
LORENZ, A. Vegetative reproduction in the New England Frullaniae, 279
Loxsomopsis, 285; costaricensis, 286; Lehmanii, 286; **notabilis,** 285, 288
Lychnis, 317, 318; affinis, 318; apetala, 319; attenuata, 319; Drummondii, 318; Kingii, 319; montana, 318; Parryi, 319; striata, 318; Taylorae, 318; triflora, 318
Lycopodiales, 232
Lysichiton camtschatcense, 107

MACKENSEN, B. Three new species of Opuntia, with a discussion of the identity of Opuntia Lindheimeri, 289
Magnolia Boulayana, 388; speciosa, 388, 390, 395
Malapoenna falcifolia, 390, 399
Malvales, 399
Marattia, 236; alata, 236; Douglasii, 236
Marattiaceae, 236
Marattiales, 232, 235
Marsilea crenulata, 233; natans, 233; quadrifolia, 233; villosa, 233
Marsileaceae, 232, 233
Matteuccia, 259
MAXON, W. R. Notes on the North American species of Phanerophlebia, 23
Maytenus **clarendonensis,** 8
Medicago lupulina, 72; sativa, 71
Meibomia canadensis, 75; obtusa, 74; rigida, 75; sessilifolia, 75

Melampsora Lini, 113, 118, 120–122, 127, 128; Rostrupi, 113, 120
Melandrium, 317, 318; affine, 318; apetalum, 319; trifolium, 318
Melanthaceae, 108
Melilotus alba, 72; officinalis, 72
Melodorum, 503; fruticosum, 503; latifolium, 504
Mertensia dichotoma, 241; emarginata, 240; excelsa, 240; glabra, 240; glauca, 239; hawaiiensis, 241; pinnata, 240
Mespilodaphne, 401
Methods of culture and the morphology of the archicarp in certain of the Ascobolaceae, 139
Mettenia **acutifolia,** 9; globosa, 9
Metzgeria, 279, 329
Metzgeriopsis, 329
Microlepia, 568, 585; brasiliensis, 585; cristata, 585; flaccida, 586; hirta, 586; japonica, 586; Khasiyana, 586; Mannii, 582; polypodioides, 586; Speluncae, 585, 586; strigosa, 585; tenuifolia, 585
Mitosis in living cells, 407
Monascus, 149, 150
Monolepis spathulata, 311
Montia Viae, 314
Mucor Mucedo, 456, 457, 464–466, 471, 473–478, 482, 484–486, 488, 489, 492–495, 497; stolonifera, 456, 457, 464–469, 471–478, 482–489, 492–497
Mucor, protoplasmic streaming in, 455
Muhlenbergia, 102; curtifolia, 102; neomexicana, 102; pauciflora, 102; polycaulis, 102; Pringlei, 102
Muscites Stolzii, 342
Myrica emarginata, 390, 393; longa, 388
Myricales, 393
Myrtales, 402

Nantucket, The ferns and flowering plants of, 69, 415
Nasturtium trachycarpum, 322
Neottopteris, 569
Nephrolepis, 568, 583; cordifolia, 583, 584; cultrifolia, 584; exaltata, 584; pendula, 583; tuberosa, 583, 584
Nephrodium cyatheoides, 599; Dubrueilianum, 599; Ecklonii, 598; exaltatum, 584; globuliferum, 591; Hudsonianum, 601; Keraudrenianum, 596; latifrons, 593; molle, 599; nymphale, 599; parasiticum, 599; polycarpon, 600; propinquum, 598; punctatum, 578; resiniferum, 598; rubiginosum, 594; squamigerum, 594; truncatum, 601; unidentatum, 597; violascens, 599
Neurogramme calomelanos, 588
New ferns from Tropical America, 285
New West Indian Lejeuneae—II, 603
Nitella, 467
Notes on the genus Widdringtonites, 341
Notes on two Jamaica plants, 12
Notes on the North American species of Phanerophlebia, 23
Notes on species of Solanum, 11
Nothochlaena ternifolia, 576

Obione phyllostegia, 312; tetraptera, 311
Observations on the degree of stomatal movement in certain plants, 15
Ocotea, 401
Octoclinis, 342
Odontoloma, 568, 582; Macraeanum, 582, 583; pulchellum, 582
Odontosoria, 568, 584; chinensis, 585
Oenothera, 19, 22, 519–521, 527, 531; biennis, 15, 21, 520, 527, 528, 532; grandiflora, 519–527, 532; Lamarckiana, 519–532; suaveolens, 520, 522, 524, 527–529, 531
Olfersia aemula, 571; gorgonea, 572
Onagra vulgaris, 527
Onoclea, 259, 273; Boryana, 378; lineata, 366; polypodioides, 271, 273; sorbifolia, 259; Spicant, 275; striata, 375
Ophioderma, 234, 235; pendulum, 235
Ophioglossaceae, 234
Ophioglossales, 233
Ophioglossum, 234; concinnum, 234, 235; ellipticum, 235; nudicaule, 235; pendulum, 235; vulgatum, 234
Opuntia **convexa,** 290; **Griffithsiana,** 291; leptocarpa, 289; Lindheimeri, 289, 290; macrorhiza, 289, 290; **reflexa,** 292; texana, 290
Opuntia Lindheimeri, Three new species of Opuntia, with a discussion of the identity of, 289
Oreodaphne, 400; **alabamensis,** 390, 400, 406
Ornithopteris aquilina, 575
Oryzopsis cuspidata, 102
Osmunda, 258, 259; borealis, 275; cinnamomea, 258; Claytoniana, 258; lineata, 366; Lunaria, 234; polypodioides, 271; procera, 383; regalis, 258; Spicant, 257, 258, 273, 275; Struthiopteris, 258, 259
Oxalidaceae, 417
Oxalis cymosa, 417; stricta, 417, 418
Oxycoccus, 551

Paesia aquilina, 575
Palaeocassia laurinea, 390, 396
Paliurus membranaceus, 401
Panicum nitidum, 102
Papilionaceae, 69
Parietaria **occidentalis,** 106; pennsylvanica, 106
Parrya macrocarpa, 326; **platycarpa,** 326
Parthenocissus, 39, 50

Pellaea, 567, 575, 579; ternifolia, 576; Weddelliana, 576
Phaeopeziza Nuttallii, 171
Phanerophlebia, 23; auriculata, 24, 26; guatemalensis, 23, 26, 28; juglandifolia, 23–25; pumila, 23, 25; macrosora, 24, 26; nobilis, 24, 25; remotispora, 23, 25; umbonata, 23, 26
Phanerophlebia, Notes on the North American species of, 23
Phegopteris crinalis, 595; Hillebrandii, 595; honolulensis, 595; Keraudreniana, 596; Keraudreniana procera, 596; microdendron, 600; punctata, 578; punctata flaccida, 579; punctata glabra, 578; sandwicensis, 598; spinulosa, 597; unidentata, 597
Phlebodium, 569
Phragmidium, 120; Potentillae-canadensis, 114, 122; speciosum, 113, 115, 120, 122; violaceum, 113, 119, 127, 128
Phycomyces nitens, 456, 457, 464, 466, 471, 488, 492, 494
Phyllites aristolochiaeformis, 388; rhomboideus, 388
Phymatodes, 569
Physaria didymocarpa lanata, 322; **lanata,** 322
Physcia parietina, 201
Picea alba, 99; albertiana, 99; canadensis, 99
PICKETT, F. L. A case of changed polarity in Spirogyra elongata, 509
Picrococcus, 549, 553; elevatus, 559; floridanus, 554, 556
Pilobolus, 170, 186
Pinaceae, 99
Pinonia, 242; splendens, 243
Pinus Mertensiana, 100
Plagiogyra, 257
Platanus primaeva, 388
Plicaria violacea, 170
Poa, 104; alpicola, 104; ampla, 104; andina, 105; arctica, 105; arida, 104, 105; brevipaniculata, 105; Buckleyana, 104, 105; caesia strictior, 105; californica, 105; callichroa, 105; crocata, 104, 105; Fendleriana, 105; flava, 104; flexuosa occidentalis, 105; Grayana, 104; idahoensis, 105; interior, 105; juncifolia, 105; Kingii, 106; laevigata, 105; laxa, 104; longiligula, 105; Multnomae, 104; nemoralis, 105; nervosa, 105; nevadensis, 105; occidentalis, 105; paddensis, 104; Pattersonii, 104; phoenicea, 104; pratensis, 104; pratericola, 105; pseudopratensis, 104; purpurascens, 104; rupicola, 105; Sandbergii, 104; serotina, 104; Sheldonii, 105; subpurpurea, 104; tenuifolia, 104; Tracyi, 105; triflora, 104
Poaceae, 102
Podophyllum, 441
Podozamites lanceolatus, 390, 391
Polyalthia, 502, 505
Polycodium, 549, 551–553, 556, 557, caesium, 555, 556; candicans, 555, 558; elevatum, 555, 558; floridanum, 555; 556, 558; **Kunthianum,** 559; Langloisii, 555, 558; melanocarpum, 555, 557, 558; neglectum, 555, 557; oblongum, 555–558; oliganthum, 555; revolutum, 555, 556; **sericeum,** 559; stamineum, 552, 555, 557, 558
Polygala cruciata, 421; polygama, 421; verticillata, 421; viridescens, 421
Polygalaceae, 421
Polygonaceae, 306
Polypodiaceae, 237
Polypodium, 285, 568; cordifolium, 583; crinale, 595; dentatum, 599; dichotomum, 240; exaltatum, 584; Filix-mas, 589; glaucum, 239; Hillebrandii, 595; honolulense, 595; **insidiosum,** 287, 288; Keraudrenianum, 596; lineare, 240; molle, 599; nymphale, 599; Paepigii, 578; parasiticum, 598; pectinatum, 239; polycarpon, 600; procerum, 596; punctatum, 578; sandwicense, 598, 600; Speluncae, 586; spinulosum, 597; stegnogrammoides, 600; trifurcatum, 285, 287; unidentatum, 597
Polystichum, 569; Dubrueilianum, 599; molle, 599; truncatum, 601
Populus, 301, 388; acuminata, 302; balsamifera, 301; balsamifera candicans, 302; Besseyana, 302; candicans, 301; Fremontii, 303; harkeriana, 390, 394; hastata, 301; Sargentii, 301, 302; trichocarpa, 301
Porella, 280, 329; platyphylla, 280
Portlandia **Harrisii,** 8; **nitens,** 10
Portulacaceae, 314
Potamogeton perfoliatus, 101; Richardsonianus, 101
Prinos padifolius, 426
Proteoides daphnogenoides, 394
Protoplasmic streaming in Mucor, 455
Pseudocarpidium avicennioides, 10; **pungens,** 10; **rigens,** 10
Pteridophyta, 232
Pteridophyta of the Hawaiian Islands, A taxonomic study of the, 227, 567
Pteridium, 567, 574; aquilinum, 575
Pteris, 567, 573, 579; alata, 574; aquilina, 575; arachnoidea, 575; arborea, 573; atropurpurea, 576; biaurita, 579; cretica, 573; decipiens, 576; decora, 577; excelsa, 573; irregularis, 573, 574; lomarioides, 574; Lydgatei, 579; melanocaulon, 574; nervosa, 573; owahuensis, 573; pedata, 576; pentaphylla, 574; peruviana, 576; psittacina, 575; regularis, 574; scab-

ripes, 574; serraria, 573; subverticillata, 576; terminalis, 573; ternifolia, 576; Treacheriana, 574; triphylla, 574
Puccinia Bunii, 117; Cirsii-lanceolati, 114–116, 124; Liliacearum, 117; Malvacearum, 128, 129; Peckiana, 116; Podophylli, 114; transformans, 114, 115
Pyronema, 63–66, 146, 147, 149, 150, 190, 191; omphalodes, 67
Pyronema, The viability of the spores of, 63

Radicula **trachycarpa,** 322
Radula, 210, 329, 338; complanata, 329, 331, 332, 336, 338; flaccida, 329–339; Hedingeri, 329, 334, 335, 337, 338, 339; Lindbergiana, 329, 338; protensa, 329, 336, 339; subtropica, 329, 338; tjibodensis, 329, 338
Radula, Discoid gemmae in, 329
Ramalina farinacea, 201; fraxinea, 201; scopulorum, 201
Ranales, 395
Ranunculaceae, 319
Ranunculus Macounii, 319; repens, 319; **rivularis,** 319
Rectolejeunea, 611; Brittoniae, 610, 611; flagelliformis, 279; **Maxonii,** 609, 611; phyllobola, 611
Relation of Ohio bog vegetation to the chemical nature of peat soils, The, 53
Rhamnales, 398
Rhamnus tenax, 390, 398
Rhizopus nigricans, 457
Rhus, 398; copallina, 423; cretacea, 397; glabra, 423; hirta, 423; reddita, 397; **reddtitiformis,** 390, 397, 406; Uddeni, 397; villosa, 398
Rhyparobius, 142, 145, 148, 149, 186, 195; albidus 143, 148; Cookei, 143, 148; crustaceus, 152; niveus, 152; pachyascus, 143, 148, 152; sexdecimsporus, 152
Robinia Pseudacacia, 74; viscosa, 74
ROBINSON, C. B. Polycodium, 549
ROBINSON, W. J. A taxonomic study of the Pteridophyta of the Hawaiian Islands, 227, 567
Rondeletia **saxicola,** 8
Rosales, 396
Ruppia curvicarpa, 101; maritima, 101
RYDBERG, P. A. Studies on the Rocky Mountain flora, 99, 301

Sabina **horizontalis,** 100; prostrata, 100
Saccobolus, 190; depauperatus, 152; Kerverni, 152; neglectus, 152, 168, 169, 193; violascens, 152
Sadleria, 569
SAFFORD, W. E. Desmos the proper generic name for the so-called Unonas of the Old World, 501
Sagittaria, 101; arifolia, 102; arifolia tenuior, 102; cuneata, 102; paniculata, 101, 102
Sagina occidentalis, 317
Salicaceae, 301
Salicales, 394
Salix, 387, 388, 439; alexensis, 304; arbusculoides, 304; arguta erythrocoma, 304; Barrettiana, 304; bella, 304; conjuncta, 304; cordata, 304; cordata angustata, 303; **cordata×monticola,** 303; deleta, 388; desertorum, 304; Drummondiana, 304; erythrocoma, 304; Fernaldii, 304; fragilis, 439; glaucops, 303; **glaucops×monticola,** 303; Hookeriana, 304; lasiolepis, 304; lucida, 304; MacCalliana, 304; monticola, 303; petiolaris, 49; **Sandbergii,** 304; saskatchewana, 304; Seemanii, 304; sitchensis, 304
Salpichlaena, 257
Salviniaceae, 232
Salviniales, 232
Sapindales, 396
Sapindus Morrisoni, 388, 390, 396
Sapranthus, 502; nicaraguensis, 502
Schizaea, 237; australis, 238; dichotoma, 237; robusta, 238
Schizostege, 568, 579; Lydgatei, 579; calocarpa, 579; pachysora, 579
Schizaeaceae, 237
SCHREINER, O., and SKINNER, J. J. The effect of guanidin on plants, 535
SEAVER, F. J. Studies in pyrophilous fungi—III. The viability of the spores of Pyronema, 63
Sequoia, 389
Seubertia, 111
Sexual fusions and spore development of the flax rust, 113
Silene, 317; Drummondii, 318
Simarubaceae, 420
Sisymbrium officinale, 323
Sitanion, 106; brevifolium, 106; ciliatum, 107; elymoides, 107; insulare, 107; lanceolatum, 107; longifolium, 106; marginatum, 107
SKINNER, J. J., and BEATTIE, J. H. Effect of asparagin on absorption and growth in wheat, 429
SKINNER, J. J., and SCHREINER, O. The effect of guanidin on plants, 535
SLOSSON, M. New ferns from Tropical America, 285
Smelowskia americana, 327; **lobata,** 327; ovata, 327
Solanum bahamense, 11; bahamense subarmatum, 11; Blodgettii, 11; boldoense, 11; verbascifolium, 11
Solanum, Notes on species of, 11
Sophia **leptostylis,** 325; procera, 325
Sordaria, 190

Sparganiaceae, 101
Sparganium angustifolium, 101; longipedunculatum, 101; simplex, 101
Sphaerotheca, 149
Spicanta attenuata, 272; borealis, 275; Ghiesbreghtii, 277; L'Herminieri, 268; onocleoides, 271; polypodioides, 272
Spirogyra communis, 509, 510; elongata, 509
Spirogyra elongata, A case of changed polarity in, 509
Sporobolus vaginaefolius, 103
Stegania, 257, 259
Stegnogramma sandwicense, 600
Stellaria alpestris, 315; borealis, 315; longipes, 315; stricta, 315; subvestita, 315; valida, 315
Stemphylium, 142
Stenanthella occidentalis, 108
Stenochlaena, 257, 372
Stenoloma tenuifolium, 585
Sterculia lugubris, 390, 399, 406
Stibasia Douglasii, 236
Stictolejeunea, 211
Stipa hymenoides, 102; membranacea, 102
Stropholirion, 111; californicum, 111
Struthiopteris, 257–259, 261, 264, 364, 373, 377; **chiriquana,** 359–362, 380; **Christii,** 360, 362, 380, 382; **costaricensis,** 360, 363, 364, 372, 384; **danaeacea,** 359, 364, 380; doodioides, 259, 275; ensiformis, 261, 262; **exaltata,** 262, 264, 266, 278; **falciformis,** 357, 365, 359, 365; **jamaicensis,** 262, 266; **L'Herminieri,** 262, 267; **lineata,** 360, 366, 368, 369, 374–377, 381–384; **Maxonii,** 262, 268; ornifolia, 373; pennsylvanica, 258; **Plumieri,** 261, 262, 266, 269, 270, 271; polypodioides, 259, 261, 270, 271, 275, 364, 383; **rufa,** 357, 358, 369; **Schiedeana,** 359, 370, 377, 380, 385; **sessilifolia,** 360, 362, 373; **Shaferi,** 357, 358, 374, 375, 378; Spicant, 258, 259, 262, 277, 278; **stolonifera,** 262, 277, 278; **striata,** 359, 360, 368, 369, 375–377, 380–385; **Underwoodiana,** 357–359, 375, 377, 380, 385; **varians,** 357, 359, 379, 380; **violacea,** 359, 373, 376, 378–382; **vivipara,** 359, 380–382; **Werckleana,** 357, 358, 364, 373, 382, 384
Struthiopteris and its representatives in North America, The genus, 257, 357
Studies in pyrophilous fungi—III. The viability of the spores of Pyronema, 63
Studies on the Rocky Mountain flora, 99, 301
Studies of West Indian plants—IV, 1
Symplocarpus foetidus, 439

Tabernaemontana discolor, 12; ochroleuca, 12
Taxonomic study of the Pteridophyta of the Hawaiian Islands, A, 227, 567
Tecoma stans, 115
Teloschistes chrysophthalmus, 201; flavicans capensis, 201
Tetraclinis, 342
Texas. Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII, 387
Tectaria, 569
Thalictrum **columbianum,** 320; megacarpum, 320; venulosum, 320
Thecotheus, 187; Pelletieri, 146, 152, 164, 166, 193
Thelebolus, 139, 186; stercoreus, 143, 146–149, 152, 167, 169, 170
Thuia graminea, 342
Thuites gramineus, 342
Thymeleales, 399
Thyopsiella, 280
Tofieldia coccinea, 108; occidentalis, 108
Toxicodendron Vernix, 424; vulgare, 424
Toxicoscordion, 110
Trachycolea, 280
Tradescantia, 107; bracteata, 108; laramiensis, 107; occidentalis, 107; scopulorum, 108; universitatis, 107; virginica occidentalis, 107
Tricalycites papyraceus, 390, 405
Trichomanes, 244; apiifolium, 245; Bauerianum, 244, 245; bifolium, 245; chinensis, 585; cuneiforme, 595; crispum, 244; cyrtotheca, 245, 246; davallioides, 246; Draytonianum, 246; eminens, 245; Endlicherianum, 246; humile, 245, 246; meifolium, 245; minutulum, 246; minutum, 245; myrioplazium, 245; parvulum, 244, 245; radicans, 245, 246; sandvicense, 246; saxifragoides, 245; sibthorpioides, 245; speciosum, 246; strigosum, 585
Trifolium arvense, 73; aureum, 72; dubium, 73; hybridum, 73; incarnatum, 73; pratense, 73, 74; procumbens, 73; repens, 74
Trillium, 439
Triphragmium, 114–116; Ulmariae, 113, 114, 122
Triteleia, 111
Tsuga, 100; Mertensiana, 100; Pattoniana, 100
Tumion, 389

Ulex europaeus, 69
Ulmaceae, 304
Umbellales, 402
Undescribed species from Cuba, 9
Undescribed species from Jamaica, 7
Unona, 501–504; acutiflora, 502; bibracteata, 501; Brandesana, 504; cochinchinensis, 506; crassipetala, 502; crinita, 505, 507; dasymaschala, 507;

desmantha, 505, 508; Desmos, 504, 506; discolor, 502, 504, 506; discreta, 502, 503; dumosa, 506; Dunalii, 506; elegans, 506; latifolia, 504; Lawii, 506; longiflora, 507; nitidissima, 502; panamensis, 501; pannosa, 506; penduliflora, 502; praecox, 507; stenopetala, 505, 507; uncinata, 502; violacea, 502; viridiflora, 506; Wrayi, 505, 507; xylopioides, 502; zeylanica, 506
Unonopsis, 501
Uromyces Caladii, 113; Lilii, 114; Scirpi, 114, 115
Urtica Breweri, 305; cardiophylla, 306; gracilis, 305, 306; **strigosissima,** 305; **viridis,** 305
Urticaceae, 305
Urticales, 394
Usnea articulata, 201; floridana, 201
Uvaria latifolia, 504; monilifera, 502
Uvularia sessilifolia, 439–441
Uvularia sessilifolia, A contribution to the life history of, 439

Vaccinium, 549–554, 556; album, 553; arboreum, 549, 551, 553; caespitosum, 552, 556; confertum, 551; consanguineum, 554, 555; corymbosum, 552; crassifolium, 553; cubense, 552; dumosum, 553; elevatum, 553, 559; frondosum, 553; Kunthianum, 552, 553, 555, 559; leucanthum, 552; melanocarpum, 559; melanocarpum candicans, 557; melanocarpum sericeum, 557–559; meridionale, 554; multiflorum, 554; myrtifolium, 553; Myrtillus, 552, 556; pallidum, 553; poasanum, 551; Ramonii, 554; scoparium, 556; stamineum, 549, 553–555, 559; stamineum melanocarpum, 557; stenophyllum, 552; Vitis-Idaea, 551, 553
Variation in the floral structures of Vitis, 37
Vatica, 406
Vegetative reproduction in the New England Frullaniae, 279
Veratrum Eschscholtzianum, 108
Viburnum robustum, 388, 390, 405
Vicia angustifolia, 79; hirsuta, 78; sativa, 79; tetrasperma, 78; villosa, 78
Viola, 92, 93; Angellae, 88–90; fimbriatula, 92; hirsutula, 95; **hirsutula** × **palmata,** 96; hirsutula × sororia, 96; **hirsutula** × **Stoneana,** 96; **hirsutula** × **triloba,** 95; latiuscula, 95; **latiuscula** × **triloba,** 94, 95; palmata, 85–89, 97; palmata dilatata, 92, 96; palmata × papilionacea, 85, 97; **palmata** × **triloba,** 88–90, 96; palmata × villosa, 95; papilionacea, 85–93, 97; **papilionacea** × **Stoneana,** 93; **papilionacea** × **triloba,** 85, 90–92; pedatifida, 97; sagittata, 92; sororia, 90, 92, 93; **sororia** × **triloba,** 92; Stoneana, 93; **Stoneana** × **triloba,** 93, 96; Stoneana × villosa, 96; triloba, 88–97; variabilis, 89, 90, 91; villosa × palmata asarifolia, 96
Violet hybrids between species of the palmata group, 85
Viscaria, 317
Vitex rigens, 10
Vitis, 37–39, 46, 50, 51; aestivalis, 43; bicolor, 38, 40, 43, 48, 49; Labrusca, 40, 48; vinifera, 39, 42, 48; vulpina, 38–45, 48, 49, 51
Vitis, Variation in the floral structures of, 37
Vitis, General floral habits of, 37; Special variation in flower parts, 42
Vittaria, 568, 586; elongata, 587; rigida, 587

Wahlbergella, 317, 318; affinis, 318; apetala, 319; **attenuata,** 319; **Drummondii,** 318; **Kingii,** 319; **montana,** 318; **Parryi,** 319; **striata,** 318; **Taylorae,** 318; triflora, 318; uniflora, 319
Was Lamarck's evening primrose (Oenothera Lamarckiana Seringe) a form of Oenothera grandiflora Solander?, 519
West Indian Lejeuneae, New—II, 603
West Indian plants, Studies of —IV, 1
West Indies, Dendropanax in the, 1
Widdringtonia, 342, 343
Widdringtonites, 342–344, 346, 347; oblongifolius, 346; Reichii, 346; subtilis, 344, 346, 347; Ungeri, 342
Widdringtonites, Notes on the genus, 341
WILLISTON, R. Discoid gemmae in Radula, 329
Woodwardia, 276

Xylopia, 504; frutescens, 504; grandiflora, 502; muricata, 504; salicifolia, 504

Zamia lanceolata, 391
Zannichelliaceae, 101
Zizyphus, 398; **lamarensis,** 390, 398, 406
Zygadenus alpinus, 108; dilatatus, 108; elegans, 108; falcatus, 110; glaberrimus, 110; gramineus, 109; intermedius, 109, 110; Nuttallii, 110; paniculatus, 110; porrifolius, 108; venenosus, 109
Zygnema pectinatum, 128